




The  
Fertility Society  
of Australia

## Pre-Conception Health Special Interest Group



### Sexually transmitted infections (STIs)

Sexually transmitted infections (STIs) are prevalent across Australia and the incidence of some STIs is rising. Some STIs can impact fertility in both men and women, affect a woman's ability to carry a pregnancy, and influence child health outcomes. The STIs that can affect fertility and/or neonatal health include chlamydia, gonorrhoea, human immunodeficiency virus (HIV), genital herpes, syphilis and *Mycoplasma genitalium*.



Your Fertility is a national public education campaign funded by the Australian Government Department of Health and Ageing under the Family Planning Grants Program.

# Sexually transmitted infections (STIs)

## Evidence review

### **Chlamydia (*Chlamydia trachomatis*)**

Chlamydia is the most commonly notified infectious disease in Australia [1] and worldwide [2]. In 2010 there were 74,305 diagnoses of chlamydia in Australia, which is an increase of 17% since 2009 [3]. Between 2003 and 2008 the rate of chlamydia diagnosis in both males and females almost doubled [1]. Infections are most common in the 15 to 29 years age-group, which accounts for 80% of cases [3]. The risk of infection is higher for women than men and in those aged 15-24 years than in those who are older [2,4]

Chlamydia is often asymptomatic so people can have the infection for some time without being diagnosed or treated [2]. In women, undetected or multiple chlamydia infections can cause pelvic inflammatory disease (PID), which in turn can lead to ectopic pregnancy and tubal infertility [2,4,5]. In men, chlamydia can affect sperm quality and function and cause urethritis, prostatitis and epididymitis [4,5] It is thought that the immune response to the infection rather than the infection itself causes the damage [4].

### **Gonorrhoea (*Neisseria gonorrhoea*)**

Gonorrhoea infection can cause similar adverse effects on fertility as chlamydia but is more commonly symptomatic. If left untreated it can lead to PID in women and infertility in both men and women. In 2009, infection rates for gonorrhoea were second only to chlamydia with over 8000 notifications [1]. Between 2009 and 2010, the number of diagnoses for gonorrhoea increased by 25% [7].

While chlamydia is more common in females, gonorrhoea is more common in males [5]. The groups most commonly affected are men who have sex with men, Indigenous Australians in remote areas and people who have had heterosexual contact overseas [5].

### **HIV (*human immunodeficiency virus*)**

HIV can be transmitted through sexual intercourse and can also be transmitted to a baby during pregnancy and through breastfeeding. HIV cannot be cured and there is currently no vaccine to prevent it, but safer sex practices can prevent transmission. With effective antiviral medications, many people are able to survive and live relatively normal lives. HIV can impair male fertility [8]. Use of assisted reproductive technology can minimise the risk of transmission in HIV discordant couples wishing to have children [10,11].

HIV is most common in men who have sex with men. In 2008, there were almost 1000 new cases of HIV in Australia [3]. Of these cases, 66% were men who have sex with men and 24% were through heterosexual contact [3].

### **Genital herpes**

Genital herpes, caused by the herpes simplex virus (HSV), is not a notifiable STI but is common among sexually active people. Genital symptoms include small, painful blisters, painful urination and vaginal discharge. Nearly 75% of women with herpes can expect at least one flare-up during pregnancy. HSV can be transmitted to the baby through direct contact with the virus during birth. To reduce the risk of transmission, it is often recommended that a cesarean be offered to women with active lesions [12].

### **Syphilis (*Treponema pallidum*)**

Syphilis can be transmitted from mother to foetus during pregnancy [1]. This is known as congenital syphilis and can result in health problems for the child at birth and later in life.

Notifications for syphilis in Australia have increased in recent years, mostly among men who have sex with men. However, notifications of congenital syphilis remain low. This is likely to be due to screening and treatment of pregnant women [1].

### ***Mycoplasma genitalium***

*Mycoplasma genitalium* is an increasingly common bacterial STI [13] which can result in vaginal discharge, urethritis, cervicitis and PID in women. PID can lead to other reproductive complications and infertility [13]. Studies have shown a strong association between *Mycoplasma genitalium* and infertility [13].

## Summary

STIs such as chlamydia, gonorrhoea, HIV, genital herpes, syphilis and *Mycoplasma genitalium* can adversely affect fertility and/or neonatal health. Notifications of some STIs continue to rise. In particular, diagnoses of chlamydia and gonorrhoea increased by 17% and 25% respectively in Australia between 2009 and 2010.

## Recommendations

Screening is an important tool in preventing the spread of chlamydial infection. Opportunistic screening may decrease infection rates and increase detection rates, allowing for earlier treatment of the infection. Bowden et al [5] recommend opportunistic screening of sexually active young women attending their doctor for other reproductive health matters such as Pap smears, contraceptive advice, antenatal care or termination of pregnancy. Transmission of STIs can be reduced or even prevented by use of safer sex practices including the use of condoms. Screening of both men and women prior to pregnancy can detect STIs and allow for treatment before pregnancy occurs. This would minimise the risk of transmission from mother to baby and prevent health problems in the unborn child.

The Department of Health and Ageing have produced helpful STI symptoms and treatment tools for health professionals [14,15].

### **Guidelines for testing of donors**

In Australia, assisted reproduction clinics are required to minimise the risk of donor gametes transmitting STIs to recipients. Guidelines produced by the National Health and Medical Research Council state that clinics should not accept gamete donations from people at increased risk of transmissible infections and that potential donors should undergo testing for such conditions [16].

The Code of Practice for reproductive technology units issued by the Reproductive Technology Accreditation Committee of the Fertility Society of Australia also requires that clinics minimise the risk of "transmission of infectious agents" between gamete/embryo donors and recipients [17].



## Sexually transmitted infections (STIs)

### References

1. Australian Institute of Health and Welfare, "Australia's Health 2010, in Australia's health series 2010." Australian Institute of Health and Welfare: Canberra.
2. Akande, V., et al., "Impact of Chlamydia trachomatis in the reproductive setting: British Fertility Society Guidelines for Practice." Human Fertility (Cambridge), 2010. 13(3): 115-125.
3. National Centre in HIV Epidemiology and Clinical Research, "HIV/AIDS, viral hepatitis and sexually transmissible infections in Australia." Annual surveillance report, NCHECR, Editor 2009, The University of New South Wales: Sydney.
4. Carey, A.J. and Beagley, K.W., "Chlamydia trachomatis, a hidden epidemic: effects on female reproduction and options for treatment." American Journal of Reproductive Immunology, 2010. 63: 576-586.
5. Bowden, F.J., et al., "Sexually transmitted infections: new diagnostic approaches and treatments." Medical Journal of Australia, 2002. 176(11): 551-557.
6. Katz, B., Fortenberry, D., and Orr, D. "Factors affecting chlamydial persistence or recurrence one and three months after treatment." In Ninth International Symposium on Human Chlamydial Infections. 1998. Napa, CA: Berkeley University Press.
7. de Wit, J., et al., "HIV/AIDS, hepatitis and sexually transmissible infections in Australia: Annual report of trends in behaviour 2011." J. de Wit, et al., Editors. 2011, National Centre in HIV Social Research: Sydney.
8. Chia SE, Ong CN, Chua LH, Ho LM, Tay SK (2000) "Comparison of zinc concentrations in blood and seminal plasma and the various sperm parameters between fertile and infertile men." Journal of Andrology 21: 53-57.
9. Ochsendorf, F.R., "Sexually transmitted infections: impact on male fertility." Andrologia, 2008. 40(2): 72-75.
10. Baker, HWG., Mijch, A., Garland, S., Crowe, S., Edgar, D., Clarke, G., Foster, P., Blood, J., "Use of assisted reproductive technology to reduce the risk of transmission of HIV in discordant couples wishing to have their own children where the male partner is seropositive with an undetectable viral load." J Med Ethics 2003, 29: 315-320.
11. Delvaux, T. and Nostlinger, C., "Reproductive choices for women and men living with HIV: contraception, abortion and fertility." Reproductive Health Matters, 2007. 15 (29 supplement): 46-66.
12. Cochrane Database of Systematic Reviews: Plain Language Summaries. <http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0013130/>
13. McGowin, C.L. and Anderson-Smits, C., "Mycoplasma genitalium: an emerging cause of sexually transmitted disease in women." PLoS Pathogens, 2011. 7(5): e1001324, [10.1371/journal.ppat.1001324](https://doi.org/10.1371/journal.ppat.1001324).
14. Department of Health and Ageing, STI Symptoms Tool  
[http://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-bbvs-sti-tools/\\$File/sti-symptoms-tool.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-bbvs-sti-tools/$File/sti-symptoms-tool.pdf)
15. Department of Health and Ageing, STI Treatment Tool  
[http://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-bbvs-sti-tools/\\$File/sti-treatment-tool.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/ohp-bbvs-sti-tools/$File/sti-treatment-tool.pdf)
16. National Health and Medical Research Council, "Ethical guidelines on the use of assisted reproductive technology in clinical practice and research." NHMRC, Editor 2007, Australian Government: Canberra.
17. Reproductive Technology Accreditation Committee. "Code of practice for assisted reproductive technology units." 2010; Available from: <http://www.fertilitysociety.com.au/rtac/>.